

The exact dynamical solution for two dust shells collapsing towards a black hole

Shuang Nan Zhang and Yuan Liu

Physics Department and Center for Astrophysics, Tsinghua University, Beijing, 100084, China¹

Abstract. The gravitational collapse of a star is an important issue both for general relativity and astrophysics, which is related to the well known “frozen star” paradox. Following the seminal work of Oppenheimer and Schneider (1939), we present the exact solution for two dust shells collapsing towards a pre-existing black hole. We find that the inner region of the shell is influenced by the property of the shell, which is contrary to the result in Newtonian theory and the clock inside the shell becomes slower as the shell collapses towards the pre-existing black hole. This result in principle may be tested experimentally if a beam of light travels across the shell. We conclude that the concept of the “frozen star” should be abandoned, since matter can indeed cross a black hole’s horizon according to the clock of an external observer. Since matter will not accumulate around the event horizon of a black hole, we predict that only gravitational wave radiation can be produced in the final stage of the merging process of two coalescing black holes. Our results also indicate that for the clock of an external observer, matter, after crossing the event horizon, will never arrive at the “singularity” (i.e. the exact center of the black hole).

Keywords: General relativity; Schwarzschild metric; Birkhoff theorem; black hole; black hole accretion; black hole merger; frozen star; black star

PACS: 04.20.-q; 04.20.Cv; 04.20.Dw; 04.20.Gz; 04.20.Jb; 04.70.-s; 04.70.Bw; 95.30.Sf; 97.60.Lf

INTRODUCTION

The “frozen star” paradox is a well known novel phenomenon predicted by general relativity, i.e. a distant observer (O) sees a test particle falling towards a black hole moving slower and slower, becoming darker and darker, and is eventually frozen near the event horizon of the black hole. However, as we discussed in [1], several possible explanations to this phenomenon have been proposed, but none of these is completely satisfactory. In the previous work, we found that an external observer should be able to observe matter falling in a black hole, based on our *stationary* solution of a dust shell around a pre-existing black hole. In this paper we further obtain the exact *dynamical* solution for two shells collapsing towards a pre-existing black hole, in particular to find the difference between this solution and that for the gravitational collapse of a uniform dust ball [2], and the implication for the “frozen star” paradox.

¹ email: zhangsn@tsinghua.edu.cn

THE DYNAMICAL SOLUTION

We follow the method adopted in the seminal work of Oppenheimer and Schneider (1939) [2], i.e. obtaining the solution in the comoving coordinates first and then transforming it into the ordinary coordinates.

The general form of the metric in the comoving coordinates is ($G = c = 1$ is adopted throughout this paper)

$$ds^2 = d\tau^2 - e^{\bar{\omega}} dR^2 - e^{\omega} (d\theta^2 + \sin^2 \theta d\phi^2), \quad (1)$$

where $\bar{\omega}(R, \tau)$ and $\omega(R, \tau)$ are the unknown functions. In the comoving coordinates, the only non-zero component of the energy momentum tensor is $T_4^4 = \rho$. As the results in [2], we can obtain the following equations from the field equation

$$e^{\omega} = (F\tau + G)^{4/3}, \quad (2)$$

$$e^{\bar{\omega}} = e^{\omega} \omega'^2 / 4, \quad (3)$$

$$8\pi\rho = \frac{4}{3} (\tau + G/F)^{-1} (\tau + G'/F')^{-1}, \quad (4)$$

where F and G are arbitrary functions of R . Dot and prime are the partial differentiation with respect to τ and R , respectively. At $\tau = 0$, we obtain

$$F = -\sqrt{6\pi\rho_0(G^2 + C_1)}, \quad (5)$$

where C_1 is the integration constant. As in [2], we choose $G = R^{3/2}$.

As obtained in [2], the value of F in region V is $-\frac{3}{2}\sqrt{r_0}$, where $r_0 = 2M$ (M is the total mass of the system). Then to assure the solutions are continuous at the boundaries, the results of F in other regions should be

$$F = \begin{cases} -\frac{3}{2}\sqrt{r_0'} & ; R < a_1' \\ -\frac{3}{2}\sqrt{\frac{8\pi\rho_1}{3}(R^3 - a_1'^3) + r_0''} & ; a_1' < R < a_1 \\ -\frac{3}{2}\sqrt{r_0''} & ; a_1 < R < a_2' \\ -\frac{3}{2}\sqrt{\frac{8\pi\rho_2}{3}(R^3 - a_2'^3) + r_0} & ; a_2' < R < a_2 \end{cases}, \quad (6)$$

where $r_0 = 2M$, $r_0' = 2m$, $r_0'' = 2(m + m_1)$.

Up to now we have obtained the solution in the comoving coordinates. Then we try to transform the solution into the ordinary coordinates, in which the metric has the form as

$$ds^2 = B(r, t) dt^2 - A(r, t) dr^2 - r^2 (d\theta^2 + \sin^2 \theta d\phi^2). \quad (7)$$

The transformation of r is obvious. We must choose $r = e^{\omega/2} = (F\tau + G)^{2/3}$.

Using the contravariant form of the metric and requiring that the g^{tr} term vanishes, we have

$$t'/\dot{t} = \dot{r}r'. \quad (8)$$




FIGURE 1. (a) is the configuration of the dynamical solution for two shells in comoving coordinates. m , m_1 and m_2 are the masses of the black hole, the shell 1 and 2, respectively. a'_1 and a_1 are the radii of the inner and outer boundaries of shell 1, respectively. a'_2 and a_2 are the radii of the inner and outer boundaries of shell 2, respectively. (b) is evolution curves of the dynamical solutions for $a_2 = 10r_0$, $a'_2 = 8r_0$, $a_1 = 5r_0$, $a'_1 = 2.5r_0$, $r'_0 = 1/3r_0$, and $r''_0 = 2/3r_0$ with coordinate time. (c) is the comparison of the evolution of the outer boundary of shell 1 between the case with (solid) or without (dashed) shell 2. The parameters of the shells are $a_2 = 10r_0$, $a'_2 = 6r_0$, $a'_1 = 2.5r_0$, $a_1 = 5r_0$, $r'_0 = 1/5r_0$, and $r''_0 = 2/5r_0$, where r_0 is the Schwarzschild radius corresponding to the total mass of the system, i.e., the sum of the masses of the pre-existing black hole, shell 1 and shell 2. The inset is the ratio of the solid line to the dashed line. It is seen clearly that outside matter, though spherically symmetric, does influence the motion of matter inside.

Substituting the expressions of r in five regions into Eq. (8), we can obtain five partial differential equations of $t(R, \tau)$.

In region V, the solution of Eq. (8) is

$$t = L(x), \quad x = \frac{2}{3\sqrt{r_0}}(R^{3/2} - r^{3/2}) - 2\sqrt{rr_0} + r_0 \ln \frac{\sqrt{r} + \sqrt{r_0}}{\sqrt{r} - \sqrt{r_0}}, \quad (9)$$

where L is an arbitrary function of x . Since the metric in the ordinary coordinates in region V is the Schwarzschild metric, $L(x)$ should be x .

In general case, the partial differential equations in other regions can be solved numerically only, using the solution in region V as the boundary condition.

After obtain the transformations in every region, we could know how the shells evolve with the coordinate time. Fig. 1a and b show the configuration of this problem and the evolution of the shells in the ordinary coordinates. In Fig. 1c, we show the evolution curves of the outer boundary of shell 1 in the case with or without shell 2. The outer boundary of shell 1 will take more time to approach the asymptotic line in the case with shell 2.

DISCUSSIONS AND CONCLUSIONS

In the Newtonian gravitation theory, if the matter is spherically symmetric, the outer matter will not influence the inner region. However, as shown in Sec. II, in general relativity case, the clock in the inner region of the shell is slower compared with the case without the shell. This effect may be testable experimentally in principle, e.g. if a beam of light travels across the shell and a parallel beam of light passes outside the shell (far

enough from the shell), then the two beams of light will travel at different velocities with respect to the observers outside the shell.

The inner shell can cross the Schwarzschild radius in the two shell case. In this sense we could observe the matter falls into a black hole and the “frozen star” paradox discussed in [3] is naturally solved. As pointed out previously by us, the origin of the “frozen star” is the “test particle” assumption for the infalling matter, which neglects the influence of the infalling matter itself on the metric [1]. In fact, for some practical astrophysical settings, the time taken for matter falling into a black hole is quite short, even for the external observer [1]. We find that the infalling matter will not accumulate outside the event horizon, and thus the quantum radiation and Gamma Ray bursts predicted in [4] and [5] are not likely to be generated. We predict that only gravitational wave radiation can be produced in the final stage of the merging process of two coalescing black holes. Future simultaneous observations by X-ray telescopes and gravitational wave telescopes shall be able to verify our prediction.

It is also interesting to note that, as can be seen from Fig. 1b, in ordinary coordinates, the matter will not collapse to the singularity ($r = 0$) even with infinite coordinate time (if $r'_0 = 0$, the inner boundary of the inner shell will take infinite time to arrive at $r = 0$). It means that in real astrophysics sense, matter can never arrive at the singularity (i.e., the exact center of the black hole) with respect to the clock of the external observer. Therefore, no gravitational singularity exists physically, even within the framework of the classical general relativity.

ACKNOWLEDGMENTS

We thank many discussions with Sumin Tang, Richard Lieu, Kinwah Wu, Kazuo Makishima, Neil Gehrels, Masaruare Shibata, Ramesh Narayan, Zheng Zhao, Zonghong Zhu and Chongming Xu. SNZ is grateful to Prof. Sandip Chakrabarti for his great effort in organizing this conference, and to the great hospitality of S.N. Bose National Centre for Basic Sciences, Kolkata, India. Many participants of this conference, especially Prof. Roy Kerr, are greatly appreciated for interesting discussions. SNZ acknowledges partial funding support by the Yangtze Endowment from the Ministry of Education at Tsinghua University, Directional Research Project of the Chinese Academy of Sciences under project No. KJCX2-YW-T03 and by the National Natural Science Foundation of China under project no. 10521001, 10733010 and 10725313.

REFERENCES

1. S.N. Zhang & Y. Liu, ASTROPHYSICS OF COMPACT OBJECTS: International Conference on Astrophysics of Compact Objects. AIP Conference Proceedings, Volume 968, pp. 384-391 [arXiv:0710.2443] (2008)
2. J.R. Oppenheimer & H. Snyder, Phys. Rev. 56, 455 (1939)
3. V.P. Frolov & I.D. Novikov, Black Hole Physics, Kluwer Academic Publishers, Dordrecht (1998)
4. T. Vachaspati, D. Stojkovic & L.M. Krauss, Phys. Rev. D 76, 024005 (2007)
5. T. Vachaspati, [arXiv:0706.1203v1] (2007)